

Efimov physics in heteronuclear four-body systems

Yujun Wang,^{1,*} W. Blake Laing,^{1,†} Javier von Stecher,² and B.D. Esry¹

¹*Department of Physics, Kansas State University, Manhattan, Kansas 66506, USA*

²*JILA, University of Colorado, 440 UCB, Boulder, Colorado, 80309, USA*

We study three- and four-body Efimov physics in a heteronuclear atomic system with three identical heavy bosonic atoms and one light atom. We show that exchange of the light atom between the heavy atoms leads to both three- and four-body features in the low-energy inelastic rate constants that trace to the Efimov effect. Further, the effective interaction generated by this exchange can provide an additional mechanism for control in ultracold experiments. Finally, we find that there is no true four-body Efimov effect — that is, no infinite number of four-body states in the absence of two- and three-body bound states — resolving a decades-long controversy.

PACS numbers:

Few-body physics has benefitted greatly from ultracold experiments on quantum gases in recent years. Many long-standing predictions have been verified and new phenomena identified due to the extraordinary ability to control and measure these systems. One of the more spectacular results for few-body physics was the observation of Efimov physics in Cs [1], experimentally validating Efimov's surprising, 35-year old prediction [2] by confirming its predicted influence on three-body recombination [3]. Several measurements of Efimov physics have now been made [1, 4–13] and even include features traceable via theory [14] to four-body processes [10, 15].

One natural question to ask is whether there is an Efimov effect for $N>3$ bodies. The Efimov effect in this case is defined as the existence of an infinite number of N -body bound states when no subsystems are bound. Part of the answer was provided in Ref. [16]: there is no Efimov effect for $N>3$ *equal mass* particles. While generally interpreted to imply no Efimov effect is possible for $N>3$ (see, for example, [17]), this result does not preclude the possibility of an Efimov effect for systems with *unequal masses*. In fact, Ref. [18] recently reported that three identical fermions interacting resonantly with a fourth particle do have an Efimov effect for a small range of mass ratios. The question of an Efimov effect for $N=4$ with three identical bosons, however, remains open despite prior study: H_3L systems (H and L are heavy and light particles, respectively) were analyzed in Ref. [19] with the conclusion that no Efimov effect occurs. When this system was revisited in Ref. [20], however, the opposite conclusion was reached.

In this Letter, we settle this controversy: there is no true four-body Efimov effect for H_3L with bosonic H s. Although our conclusion is in agreement with Ref. [19], our reasoning is very different. We have, however, identified one universal four-body state attached to each three-body Efimov threshold. We also show that the low-energy scattering observables simultaneously display distinct three- and four-body features characteristic of the three-body Efimov effect. Moreover, we find that, in the context of ultracold collisions, the s -wave two-body scat-

tering length a_{HL} between H and L atoms can be used to tune the effective heavy-heavy scattering length a_{HH}^* when $a_{HL} > 0$, opening up new avenues for control in few-body systems.

Since a direct solution of the four-body problem remains a substantial challenge, we apply the Born-Oppenheimer (BO) approximation as described in Refs. [19, 20], assuming that the mass m of L is much less than the mass M of H . Although we will focus on bosonic H s, our analysis can be generalized straightforwardly.

We first recall that the three-body Efimov effect occurs when at least two scattering lengths are much larger than a characteristic two-body interaction range r_0 [2]. Thus, for H_2L , Efimov physics will occur for $|a_{HL}| \gg r_0$. Applying the BO approximation with ρ the $H+H$ distance yields a BO potential with the universal long-range behavior $-\chi_0^2 \hbar^2 / 2m\rho^2$, $\chi_0 \approx 0.567143$, in the region $r_0 \ll \rho \ll |a_{HL}|$ [17, 21]. The three-body problem has thus been reduced to an effective two-body problem whose interaction is the Efimov potential, giving the characteristic geometric relation between bound state energies

$$E_{n+1}/E_n = e^{-2\pi/s_0}, \quad n = 0, 1, 2, \dots \quad (1)$$

where $s_0^2 = \chi_0^2 M / 2m - 1/4$. An infinite number of Efimov states are thus obtained in the limit $|a_{HL}| \rightarrow \infty$, corresponding to a zero-energy HL bound state.

Similarly, all of the universal results [17] for low-energy three-body scattering apply to H_2L . For instance, when $a_{HL} > 0$, there is a weakly bound HL molecule, and

$$a_{HH}^* \propto \cot[s_0 \ln(a_{HL}/r_0) + \Phi] a_{HL} \quad (2)$$

where r_0 is the characteristic range of the two-body interaction. The poles in a_{HH}^* occur at those values of a_{HL} when an Efimov state becomes bound, so their positions are characteristically log-periodic in a_{HL} . Their overall position, however, is determined by the short-range three-body phase Φ [3, 22]. Equation (2) thus shows that a_{HL} provides a control over a_{HH}^* that could prove advantageous when a_{HH}^* cannot easily be controlled directly via, for example, a Feshbach resonance [23].

In general, the H s also interact directly. However, so long as the direct $H+H$ interaction is short-ranged — even if it is repulsive — the large- ρ behavior of the effective $H+H$ interaction is unchanged, and the Efimov effect remains. A direct $H+H$ interaction changes only E_0 (or, equivalently, Φ) and not the characteristic log-periodic behavior. Consequently, manipulating the direct $H+H$ interaction allows control of Φ and thus the position of the family of log-periodic Efimov features.

Applying the BO approximation to H_3L reduces it to an effective three-body problem for H_3 . We thus expect that our knowledge of Efimov physics and universality for three identical bosons should apply and provide at least a basic understanding of the system. For simplicity, we assume that the H atoms do not interact directly. Per the argument above, we know this will not materially affect our conclusions. The $H+H$ interaction then comes solely from mediation by the L and is characterized by the effective $H+H$ scattering length a_{HH}^* from Eq. (2). Based on the result for three bosons, we expect an Efimov effect for H_3 when $|a_{HH}^*| \rightarrow \infty$ [20].

To determine whether the $|a_{HH}^*| \rightarrow \infty$ limit actually produces an Efimov effect for H_3L , we adopt the following definition [16, 19]: a true four-body Efimov effect exists if the four-body system possesses an infinity of stable bound states when there is a zero-energy three-body bound state and no other two- or three-body bound states. Under these conditions, the four-body Efimov effect is completely independent of any three-body Efimov effect, making the resulting states fundamentally different from the universal four-body states discussed in Refs. [14, 24–26].

If it exists, a true four-body H_3L Efimov effect will occur for $|a_{HH}^*| \rightarrow \infty$ and $a_{HL} < 0$. This case gives a zero-energy H_2L bound state but no HL bound state and is precisely the case identified in Ref. [19]. They concluded that no Efimov effect is possible because the BO potential surface has no long-range component. Naus and Tjon [20] correctly pointed out that it is not the BO potential surface that must behave as $-R^{-2}$, but rather the adiabatic hyperspherical potential where the hyperradius R measures the overall size of the system (see, for example, Refs. [27, 28] for a discussion of hyperspherical coordinates in this context). They concluded that because $|a_{HH}^*| \rightarrow \infty$, the conditions of the Efimov effect for H_3 are fulfilled and there is thus a four-body Efimov effect. Unfortunately, neither analysis provides a definitive answer.

Although otherwise sound, the argument of Ref. [20] is limited by its reliance on the BO approximation — a problem they identified but did not address. The issue with the BO approximation is that for $a_{HL} < 0$, L becomes unbound when $\rho \gtrsim |a_{HL}|$ [19], and its spectrum becomes continuous. Since a zero-energy L is no longer fast compared to the H s, the BO approximation breaks down. Naus and Tjon followed Ref. [19] and simply set the BO




FIG. 1: Schematic energy spectrum for H_3L . Solid lines denote bound states and dashed lines denote resonances.

potential at such points to zero. Under this questionable approximation, a_{HH}^* can still be defined and adjusted to infinity, so that they conclude there is an infinite series of bound H_3L states, i.e. the four-body Efimov effect.

Where the BO approximation fails, however, the adiabatic hyperspherical approximation is perfectly valid. In particular, the lowest H_2L adiabatic hyperspherical potential, which corresponds to the BO potential, crosses zero energy at a hyperradius $R \sim |a_{HL}|$ much like the BO potential, but remains discrete and increases to a barrier with height $\sim 1/a_{HL}^2$ [29]. For R beyond this barrier, the potential falls to zero as $l_{\text{eff}}(l_{\text{eff}} + 1)\hbar^2/MR^2$ with $l_{\text{eff}} = 3/2$ [30].

Having taken care to define the H_2L hyperradius R such that it reduces to the $H+H$ distance ρ in the $M/m \rightarrow \infty$ limit, we can regard this adiabatic hyperspherical potential as an effective $H+H$ interaction. Since the Efimov effect for the three H s can only occur for s -wave two-body interactions, i.e. $l_{\text{eff}} = 0$, the fact that $l_{\text{eff}} = 3/2$ prohibits the Efimov effect. This conclusion has been confirmed by direct calculations of the four-body adiabatic hyperspherical potentials for H_3L with $M/m = 30$ [31] using the correlated Gaussian approach [32].

Even though our finding no true four-body Efimov effect agrees with the conclusions of Ref. [19] and contradicts the conclusions of Ref. [20], we believe the former were right for the wrong reason and the latter underestimated the consequences of the breakdown of the BO approximation. In the end, it is this breakdown that excludes the possibility of a true four-body Efimov effect in this system.

The possibility of true four-body Efimov states is not the only phenomenon of interest in H_3L . Continuing to $a_{HL} > 0$ such that $a_{HL} \gg r_0$, the BO approximation displays no pathologies since L is bound for all H configurations, and we can safely think about the three-body H_3 motion on the lowest BO potential surface. Based on

the known three-body results [17, 27, 33], we expect an adiabatic hyperspherical potential of the form

$$W_0 = -\frac{g_0^2 + 1/4}{2\mu R^2} \hbar^2, \quad a_{HL} \ll R \ll |a_{HH}^*|, \quad (3)$$

with $g_0=1.00624$ and $\mu=M/\sqrt{3}$ that approaches the three-body break-up threshold. In this case, that threshold corresponds physically to $HL+H+H$. The lower limit of R in Eq. (3) is modified from the usual three-body problem [33] due to the fact that the characteristic range of the effective two-body $H+H$ potential is no longer r_0 , but rather a_{HL} as defined by the size of the HL bound state.

When a_{HL} is tuned to give $a_{HH}^* \rightarrow \infty$, the Efimov potential (3) extends to infinity, producing an infinite series of four-body bound states below the $HL+H+H$ threshold with binding energies $E_{n+1}/E_n = e^{-2\pi/g_0}$. These states are not true four-body Efimov states since there is an HL bound state, but they can be regarded as three-body Efimov states of $HL+H+H$. They are indicated with the notation $(HL)H_2$ in Fig. 1 where the H_2L+H and $HL+H+H$ thresholds intersect since $a_{HH}^* \rightarrow \infty$ at these points.

Figure 1 sketches the energy trajectories for H_3L as a function of a_{HL} . In addition to the $(HL)H_2$ Efimov states, there are the H_2L Efimov states, and associated with each of these we find — for both $M/m=50$ and $M/m=30$ — one H_3L state that appears to be the analog of the universal four-boson states in Refs. [14, 15, 24–26]. We find that its binding energy is universally related to the binding energy of the associated H_2L Efimov state by $E_{H_3L}/E_{H_2L} \approx 0.4$ for both of the mass ratios we have calculated. Note that the binding energies are defined relative to the next lowest breakup threshold ($HL+H+H$ for H_2L states and H_2L+H for H_3L states).

Figure 2 shows our numerically calculated three-body recombination rates K_3 for $HL+H+H \rightarrow H_2L+H$ with $M/m=30$ (see Ref. [28] for details of our numerical methods). Although there may be several final H_2L Efimov states available, our calculation shows that recombination into the most weakly bound Efimov state dominates. In fact, the main peaks in Fig. 2(c) occur where an H_2L Efimov state just becomes bound. The separation between the main peaks is thus determined by s_0 from the H_2L Efimov effect: for $M/m=30$, $a_{HL}^{(2)}/a_{HL}^{(1)} = e^{\pi/s_0} = 4.34$. Note that the adiabatic hyperspherical approach gives $e^{\pi/s_0} = 3.96$ [33, 34], giving an indication of the BO approximation error for this mass ratio. For consistency, we will quote only BO results in the rest of this Letter. The factor of 5.5 between these two main peaks in Fig. 2 does not match this prediction because the criterion $a_{HL} \gg r_0$ is not well satisfied.

Since each main peak corresponds to a pole of a_{HH}^* , K_3 shows Efimov features characteristic of the H_3 motion where $|a_{HH}^*| \gg a_{HL}$. When this condition is satisfied,




FIG. 2: K_3 for $HL+H+H \rightarrow H_2L+H$ when $M/m=30$. The main peaks in (c) reflect the Efimov physics of H_2L and occur where (d) a_{HH}^* diverges. The expanded plots (a) and (b) highlight the features on each main peak related to H_3L states. The vertical dashed lines indicate where $a_{HH}^* = \infty$.

the rates are given by the usual universal three-body expressions [17, 35] with the short-range length scale set to a_{HL}

$$K_3^{(a_{HH}^* < 0)} = \frac{C'}{\mu} \frac{\sinh(2\eta) |a_{HH}^*|^4}{\sin^2[g_0 \ln(|a_{HH}^*|/a_{HL}) + \Phi] + \sinh^2(\eta)}$$

$$K_3^{(a_{HH}^* > 0)} = \frac{C}{\mu} \sin^2[g_0 \ln(a_{HH}^*/a_{HL}) + \Phi] (a_{HH}^*)^4. \quad (4)$$

In these expressions, C and C' are universal constants. But, because the final H_2L state is an Efimov state, Φ , Φ' and η depend not on short-range four-body physics, but rather on the short-range physics of the H_2L states — no additional four-body parameter is needed [14, 24, 25]. The $a_{HH}^*-K_3$ projection in Fig. 3 supports this conclusion, showing that the positions of the Efimov features related to the H_3 motion described by Eq. (4) are approaching a universal position in the limit $a_{HL} \gg r_0$.

Interestingly, for larger mass ratios, H_2L Efimov states with non-zero orbital angular momentum j are possible [33, 34]. In this case, the universal constant s_0 in Eq. (1) is determined from $s_0^2 = \chi_0^2 M/2m - j(j+1) - 1/4$, and there will be an Efimov effect for H_2L so long as $s_0^2 > 0$. Higher angular momentum Efimov states have not yet been observed because experiments have focused mostly on identical particles which have no such states and because $j > 0$ Efimov states produce extremely narrow recombination peaks as seen in Fig. 3. The mass ratio in Fig. 3, $M/m=50$, supports $j=2$ Efimov states, but is not sufficient for higher j states. Because $l_{\text{eff}} = j \neq 0$ for the H_2L system, there is no $(HL)H_2$ Efimov effect associated with these states. Their narrow K_3 peaks thus show no substructure of the sort seen on the main peaks.

The periods for each of the three different families of Efimov peaks are indicated in Fig. 3. Because none of the




FIG. 3: K_3 for $HL+H+H \rightarrow H_2L+H$ with $M/m=50$ simultaneously showing the dependence on a_{HL} and on a_{HH}^* . The H_2L Efimov physics is highlighted by the a_{HL} - K_3 projection while the a_{HH}^* - K_3 projection illustrates the $(HL)H_2$ Efimov physics. Note that a_{HH}^* increases into the figure.

respective scattering lengths are strongly in the universal limit, however, the calculated periods do not match the predicted ones. The main $j=0$ H_2L Efimov peaks highlighted by the a_{HL} - K_3 projection should have a period of $e^{\pi/s_0}=3.08$. The calculated spacings are larger than this, but appear to be approaching the expected value as a_{HL} increases. Similarly, for $j=2$, the expected period is 10.46 while the calculated one is 17.5, and the predicted period of the $(HL)H_2$ Efimov substructure on the main peaks highlighted in the a_{HH}^* - K_3 projection is $e^{\pi/g_0}=22.7$. The magnitude of the deviations from the predictions likely reflects what can be observed since it is difficult to penetrate deeply into the universal regime experimentally.

So far, we have not taken advantage of all of the freedom that this heteronuclear system affords to manipulate the Efimov features. In particular, since a_{HH}^* depends only on the total $H+H$ interaction, i.e. effective plus direct interactions, it can be tuned by either interaction — or both. Intriguingly, this freedom also allows experimental control of both a_{HH}^* and Φ (or Φ'). Controlling the former via either a_{HL} or a direct interaction with scattering length a_{HH} allows the various Efimov features seen, for instance, in Figs. 2 and 3 to be mapped out. Controlling the latter makes it possible to shift all of these features — something not possible so far in three-body systems. Moreover, in the neighborhood of a pole in a_{HH}^* , it should be possible to exert both types of control largely independently.

To illustrate the effect of tuning the direct interaction, we show in Fig. 4 the numerically calculated rates for the relaxation process $H_2L(n)+H \rightarrow H_2L(n-1)+H$ as a function of a_{HH} where n labels the most weakly bound H_2L state. This tuning was accomplished by including a short-range, direct $H+H$ interaction in addition to an




FIG. 4: V_{rel} for $H_2L(n)+H \rightarrow H_2L(n-1)+H$ for (a) $n=1$ and (b) $n=2$. The dashed lines indicate the positions where $|a_{HH}^*|=\infty$.

effective $H+H$ interaction. The behavior of V_{rel} in Fig. 4 is also found in the three-boson system. In fact, when $a_{HH}^* \gg a_{HL} > 0$, V_{rel} has the same form as in three-boson systems [17]

$$V_{\text{rel}}^{(a_{HH}^* > 0)} = \frac{A \sinh 2\eta}{\sin^2[g_0 \ln(a_{HH}^*/a_{HL}) + \Phi] + \sinh^2 \eta} a_{HH}^*, \quad (5)$$

where A is a universal constant, but η and Φ depend on the short-range details of the relaxed H_2L bound state — although not on a separate four-body parameter.

To conclude, we have studied Efimov physics in the four-body heteronuclear system H_3L with bosonic H atoms, showing that there is no true four-body Efimov effect. We have, however, identified a universal four-body state linked to each H_2L Efimov state. Perhaps more significantly, we have shown that the richness of the heteronuclear system can be exploited to reveal a variety of Efimov features with different universal scaling by scanning just a_{HL} . That same richness allows more opportunities for controlling the system as well, which is potentially very important for experiments seeking to study or utilize these processes. For instance, taking advantage of the BO approach introduced in Ref. [19], we pointed out that scanning a_{HL} gives a means for controlling the scattering length between the heavy atoms — a tool that could prove useful experimentally. Heteronuclear four-body and larger systems thus provide considerable possibilities for studying Efimov physics that were not previously anticipated.

We thank C.H. Greene for critical and stimulating discussions of the four-body Efimov effect. This work was supported in part by the National Science Foundation and in part by the Air Force Office of Scientific Research. Y. W. also acknowledges support from the National Science Foundation under Grant No. PHY0970114.

* Present address: JILA, University of Colorado, 440 UCB, Boulder, Colorado, 80309, USA

† Present address: Department of Physics and Astronomy, Rowan University, Glassboro, New Jersey 08028, USA

- [1] T. Kraemer, *et al.*, Nature **440**, 315 (2006).
- [2] V. Efimov, Phys. Lett. B **33**, 563 (1970).
- [3] B. D. Esry, *et al.*, Phys. Rev. Lett. **83**, 1751 (1999).
- [4] S. Knoop, *et al.*, Nat. Phys. **5**, 227 (2009).
- [5] T. B. Ottenstein, *et al.*, Phys. Rev. Lett. **101**, 203202 (2008).
- [6] J. H. Huckans, *et al.*, Phys. Rev. Lett. **102**, 165302 (2009).
- [7] G. Barontini, *et al.*, Phys. Rev. Lett. **103**, 043201 (2009).
- [8] M. Zaccanti, *et al.*, Nat. Phys. **5**, 586 (2009).
- [9] N. Gross, *et al.*, Phys. Rev. Lett. **103**, 163202 (2009); *ibid.*, **105**, 103203 (2010).
- [10] S. E. Pollack, *et al.*, Science **326**, 1683 (2009).
- [11] T. Lompe, *et al.*, Science **330**, 940 (2010).
- [12] M. Berninger, *et al.*, arXiv:1106.3933 (2011).
- [13] J. R. Williams, *et al.*, Phys. Rev. Lett. **103**, 130404 (2009).
- [14] J. von Stecher, *et al.*, Nat. Phys. **5**, 417 (2009).
- [15] F. Ferlaino, *et al.*, Phys. Rev. Lett. **102**, 140401 (2009).
- [16] R. D. Amado and F. C. Greenwood, Phys. Rev. D **7**, 2517 (1973).
- [17] E. Braaten and H. W. Hammer, Phys. Rep. **428**, 259 (2006).
- [18] Y. Castin, *et al.*, Phys. Rev. Lett. **105**, 223201 (2010).
- [19] S. K. Adhikari and A. C. Fonseca, Phys. Rev. D **24**, 416 (1981).
- [20] H. Naus and J. A. Tjon, Few-Body Syst. **2**, 121 (1987).
- [21] A. C. Fonseca, *et al.*, Nucl. Phys. A **320**, 273 (1979).
- [22] J. P. D’Incao, *et al.*, J. Phys. B **42**, 044016 (2009).
- [23] C. Chin, *et al.*, Rev. Mod. Phys. **82**, 1225 (2010).
- [24] L. Platter, *et al.*, Phys. Rev. A **70**, 052101 (2004).
- [25] H. W. Hammer and L. Platter, Eur. Phys. J. A **32**, 113 (2007).
- [26] R. Schmidt and S. Moroz, Phys. Rev. A **81**, 052709 (2010).
- [27] E. Nielsen, *et al.*, Phys. Rep. **347**, 373 (2001).
- [28] H. Suno, *et al.*, Phys. Rev. A **65**, 042725 (2002).
- [29] B. D. Esry, *et al.*, J. Phys. B **29**, L51 (1996).
- [30] B. D. Esry, *et al.*, Phys. Rev. A **65**, 010705 (2001).
- [31] J. von Stecher, unpublished.
- [32] J. von Stecher and C. H. Greene, Phys. Rev. A **80**, 022504 (2009).
- [33] V. Efimov, Nucl. Phys. A **210**, 157 (1973).
- [34] J. P. D’Incao and B. D. Esry, Phys. Rev. A **73**, 030702 (2006).
- [35] J.P. D’Incao and B.D. Esry, Phys. Rev. Lett. **94**, 213201 (2005).